



SCHOLARLY PUBLICATIONS
School of Biotechnology
KIIT Deemed to be University

Journal Name: Molecular Cancer


IF: 33.9

Title: Epithelial-to-mesenchymal transition (EMT) and cancer metastasis: the status quo of methods and experimental models 2025

Author: Allgayer H.; Mahapatra S.; Mishra B.; Swain B.; Saha S.; Khanra S.; Kumari K.; Panda V.K.; Malhotra D.; Patil N.S.; Leupold J.H.; Kundu G.C.

Details: Volume 24, Issue 1, December 2025, Article number 167

Abstract: Epithelial-to-mesenchymal transition (EMT) is a crucial cellular process for embryogenesis, wound healing, and cancer progression. It involves a shift in cell interactions, leading to the detachment of epithelial cells and activation of gene programs promoting a mesenchymal state. EMT plays a significant role in cancer metastasis triggering tumor initiation and stemness, and activates metastatic cascades resulting in resistance to therapy. Moreover, reversal of EMT contributes to the formation of metastatic lesions. Metastasis still needs to be better understood functionally in its major but complex steps of migration, invasion, intravasation, dissemination, which contributes to the establishment of minimal residual disease (MRD), extravasation, and successful seeding and growth of metastatic lesions at microenvironmentally heterogeneous sites. Therefore, the current review article intends to present, and discuss comprehensively, the status quo of experimental models able to investigate EMT and metastasis in vitro and in vivo, for researchers planning to enter the field. We emphasize various methods to understand EMT function and the major steps of metastasis, including diverse migration, invasion and matrix degradation assays, microfluidics, 3D co-culture models, spheroids, organoids, or latest spatial and imaging methods to analyze complex compartments. In vivo models such as the chorionallantoic membrane (CAM) assay, cell line-derived and patient-derived xenografts, syngeneic, genetically modified, and humanized mice, are presented as a promising arsenal of tools to analyze intravasation, site specific metastasis, and treatment response.



URL: <https://molecular-cancer.biomedcentral.com/articles/10.1186/s12943-025-02338-2>





SCHOLARLY PUBLICATIONS School of Biotechnology KIIT Deemed to be University

Journal Name: Renewable Energy


IF: 9.0

Title: Biodiesel synthesis from *Ricinus communis* and *Pongamia pinnata* oil blends by injecting superheated methanol & isopropanol mixtures: Optimization through CCD and ANN approaches

Author: Karmakar B.; Chakraborty S.; Kumar R.; Halder G.

Details: Volume 249, 15 August 2025, 123223

Abstract: In the current study, blends of castor and karanja oils were subjected to uncatalysed alcoholysis with superheated mixtures of 2-propanol and methanol for their rapid conversion into fuel-grade esters. Optimizable ranges identified from batch studies for 6 parameters: alcohol preheat temperature, castor oil to karanja oil ratio, initial oil mass, methanol to 2-propanol ratio, reaction temperature and retention duration were fed into a spherical central composite design (CCD-S, used for identifying process conditions for optimal biodiesel yield. It was noted that a maximum biodiesel yield of 98.79 % could be obtained when 650g castor and karanja oil blend at a ratio of 2:1 was charged into the reactor. The alcohols at a ratio of 3:5 for methanol: 2-propanol had to be pre-heated to 140 °C to achieve desired energy, reactivity and flow. The reaction provided best results when allowed to occur at 260 °C for a duration of 8 min. The experimentally obtained data were verified for reliability through ANOVA studies and ANN was used to validate the data as well as develop a model capable of predicting output accurately, with a 6-10-1 algorithm giving an R^2 of 0.987, indicating high reliability.



URL: <https://www.sciencedirect.com/science/article/pii/S0960148125008857?via%3Dihub>





SCHOLARLY PUBLICATIONS School of Biotechnology KIIT Deemed to be University

Journal Name: Plant Stress


IF: 6.8

Title: Pearl millet WRKY transcription factor PgWRKY52 positively regulates salt stress tolerance through ABA-MeJA mediated transcriptional regulation

Author: Chanwala J.; Kumari K.; Jha D.K.; Giri M.K.; Dey N.

Details: Volume 16, June 2025, Article number 100814

Abstract: Environmental stresses adversely affect plant growth and development by disturbing physiological and metabolic equilibrium. Plants counteract these stresses through intricate genetic and biochemical pathways, which are largely mediated by signalling networks that involve key transcription factors (TFs). Among these, WRKY TFs are crucial in modulating plant responses to various stresses. In previous studies, WRKY TFs have been identified in millets. However, their functional characterization in millets remains vastly unexplored. Therefore, we have isolated and characterized PgWRKY52, a Group IIc WRKY TFs from pearl millet, along with its upstream promoter region to understand its functional regulatory role. Ectopic expression of PgWRKY52 in transgenic Arabidopsis improved seed germination under salt stress and phytohormonal treatments of abscisic acid (ABA) and methyl jasmonate (MeJA). Improved stress tolerance was linked to reduced reactive oxygen species (ROS) accumulation and upregulation of stress-responsive genes, indicating an enhanced defense system. Promoter analysis unveiled that the PgWRKY52 promoter was constitutively active across vegetative and reproductive tissues, with strong stress-inducible activity under salt, heat, and ABA treatments. Cis-regulatory element (CRE) analysis identified key stress-responsive elements, including ABRE, MYB, W-box and MYC, which were validated through mutational studies as essential for promoter activity. Additionally, PgWRKY52 exhibited W-box-dependent DNA-binding capability, a characteristic feature of WRKY TFs. These findings emphasize the important function of the PgWRKY52 promoter in driving stress-responsive transcription. Altogether, these findings establish PgWRKY52 as a stress-responsive TF that enhances salt stress tolerance through crosstalk of ABA-MeJA signalling pathways and the regulatory role of its promoter, presenting a promising tool for developing climate-resilient crops.



URL: <https://www.sciencedirect.com/science/article/pii/S2667064X2500079X?via%3Dihub>





SCHOLARLY PUBLICATIONS

School of Biotechnology

KIIT Deemed to be University

Journal Name: LWT

IF: 6.6

Title: Formulation and characterization of ready-to-drink nutraceutical beverage from blood fruit (*Haematocarpus validus*)

Author: Sasikumar R.; Kumar T S.; Vivek K.; Panda S.K.; Jaiswal A.K.

Details: Volume 2251, June 2025, Article number 117929

Abstract: Blood fruit (*Haematocarpus validus*) is a rich source of bioactive compounds with potential health benefits. This study aimed to develop and characterize a ready-to-drink (RTD) fermented nutraceutical beverage (FNB) using controlled fermentation at 20, 23, and 26 °C and to assess its biochemical composition, antioxidant potential, volatile profile, anti-nutritional factors, and sensory attributes. The beverage was analyzed using Fourier Transform Infrared (FTIR), Ultra Performance Liquid Chromatography (UPLC), High Performance Liquid Chromatography (HPLC), Gas Chromatography High Resolution Mass Spectrometry (GC-HRMS), and High-Resolution Liquid Chromatography with Mass Spectrometry (HR-LCMS), with key parameters such as total sugars, organic acids, total polyphenols, flavonoids, anthocyanins, and antioxidant activity determined. Fermentation at 26 °C resulted in the highest enrichment of total anthocyanins (411.7 ± 0.79 mg C3GE/100 mL), flavonoids (481.8 ± 0.49 mg RE/100 mL), and polyphenols (537.4 ± 0.84 mg GAE/100 mL). Antioxidant activity was significantly higher in the 26 °C fermented beverage, as indicated by 2,2-diphenyl-1-picrylhydrazyl or DPPH (2128 ± 2 μ mol TE/100 mL), Oxygen Radical Absorbance Capacity or ORAC (2988 ± 3 μ mol TE/100 mL), 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid or ABTS (1238 ± 2 μ mol TE/100 mL), and Ferric Reducing Antioxidant Power or FRAP (3360 ± 2 μ mol Fe²⁺/100 mL). Volatile analysis identified 36 key compounds, contributing to sensory attributes.



URL: <https://www.sciencedirect.com/science/article/pii/S0023643825006139?via%3Dihub>





SCHOLARLY PUBLICATIONS
School of Biotechnology
KIIT Deemed to be University

Journal Name: Journal of Industrial and Engineering Chemistry


IF: 5.9

Title: A multi-approach study on CO₂ absorption in packed beds: Theoretical, experimental, and CFD perspectives on gas phase pulsation

Author: Pattnaik C.; Kumar R.; Khan M.A.; Pahari P.; Banik A.; Jeon B.-H.; Banerjee S.; Chakraborty S.; Tripathy S.K.

Details: Volume 146, Pages 641 – 655, 25 June 2025

Abstract: This work seeks to improve CO₂ absorption efficiency in packed bed columns by substituting amine-based solvents with sodium hydroxide and implementing gas phase pulsation to enhance mass transfer coefficients. Experimental analysis and computational fluid dynamics modeling were employed to investigate the impact of pulsation on absorption efficiency under various conditions. Essential parameters comprised superficial liquid velocity (1.2–4.6 cm/s), pulsation frequency (0–10 Hz), amplitude (0–20 mm), and NaOH concentration (0.25 N to 2 N), while maintaining a constant superficial gas velocity of 120 cm/s and a solute gas concentration of 13 %. Three packing materials—glass spheres, ceramic Raschig rings, and ceramic Pall rings—were evaluated. The results demonstrated that ceramic Pall rings exhibited the greatest efficiency. Pulsation, namely at 9.06 Hz and 20 mm amplitude, enhanced the volumetric mass transfer coefficient by as much as 4.53 times for Pall rings. Increased column diameters (from 7.00 cm to 11.5 cm) enhanced performance. The findings show advancement of more efficient CO₂ absorption (by switching from chemical absorption using amine based solvents to classical chemical absorption using aqueous NaOH solution) for industrial applications, aiding climate change mitigation initiatives.



URL: <https://www.sciencedirect.com/science/article/pii/S1226086X24007858?via%3Dihub>





SCHOLARLY PUBLICATIONS School of Biotechnology KIIT Deemed to be University

Journal Name: Biomass and Bioenergy

IF: 5.8

Title: Fermentation of sugarcane bagasse for production of value-added phenolic compounds using potential bacterial strains: A comparative analysis

Author: Pattnaik B.; Preeti; Gupta D.; Deb D.; Selvaraj M.; Assiri M.A.; Mohapatra S.R.; Sahoo H.P.; Tapas S.; Sarangi P.K.

Details: Volume 202, November 2025

Abstract: The present study investigates the potential of bacterial strains, viz., *Pseudomonas fragi*, *Lactobacillus plantarum*, and *Lactobacillus acidophilus*, for the production of phenolic compounds from sugarcane bagasse (SCB). The important bio-transformed phenolic products isolated from the medium were ferulic acid (FA), vanillin and vanillic acid (VA), whose identification and quantification were done by high-performance thin-layer chromatography. Carbohydrate concentration from the de-starched bagasse was also assessed and compared with that of the original (control) bagasse. Results revealed that the utmost FA yield per kg of SCB was 275 mg from *Lactobacillus acidophilus*, 225 mg from *Pseudomonas fragi* on the 9th day, and 212 mg from *Lactobacillus plantarum* on the 12th day of incubation. Likewise, the peak vanillin and VA quantified per ml of fermented extract were 16 mg on 9th and 12th day of incubation, respectively, for *Lactobacillus plantarum*, 14 mg of vanillin and 13 mg of VA on 9th day for *Pseudomonas fragi*. However, in *Lactobacillus acidophilus* 15 mg of Vanillin and 18 mg of VA was recorded on 12th day of incubation. To compare enzymatic efficiency and structural integrity among ferulic acid esterases (FAEs), a 3D structural model was constructed. We first time demonstrated that the lid domain's structural integrity enhances enzyme efficiency which has been expressed in terms of yield. An ~18 % higher yield of primary phenolic compound was obtained for *L. acidophilus* with compact FAE lid domain compared to PsfFAE. This finding highlights the metabolic potential of these strains for phenolics production and their relevance in biotransformation processes.



URL: <https://www.sciencedirect.com/science/article/abs/pii/S0961953425005720?via%3Dihub>





SCHOLARLY PUBLICATIONS School of Biotechnology KIIT Deemed to be University

Journal Name: Frontiers in Immunology


IF: 5.7

Title: Designing a potent multivalent epitope vaccine candidate against *Orientia tsutsugamushi* via reverse vaccinology technique - bioinformatics and immunoinformatic approach

Author: Panda S.; Swain S.K.; Sahu B.P.; Mahapatra S.R.; Dey J.; Sarangi R.; Ranade A.V.; Mishra N.

Details: Volume 16, 2025, Article number 1513245

Abstract: Scrub typhus is a life-threatening, undifferentiated febrile illness caused by a gram-negative bacterium, *Orientia tsutsugamushi*. The bacterial strain is a global health concern that should be considered. Despite several years of effort for the development of an effective immunogenic vaccine, no successful licensed vaccine is available. The aim of the study is to construct an epitope response using a reverse vaccinology approach. The TSA56 and ScaA proteins combined can be the most promising subunit vaccine candidates against *O. tsutsugamushi*. B-cell, CTL, and HTL epitopes were predicted, and subsequently, all the epitopes were linked by KK, AAY, and GP GPG linkers, respectively, along with an adjuvant at the N-terminal region. Furthermore, molecular docking and MD simulations were performed that exhibited a higher affinity towards TLR-2. A total of 16 linear B-cells, 6 CTL, and 2 HTL epitopes were identified and validated. The final vaccine construct showed high antigenicity, stability, and solubility. Molecular docking and MD simulations indicated strong binding interactions with TLR-2 and a stable vaccine-receptor complex. The expression of the vaccine in pET28a (+) vector was successfully implemented via in silico cloning as well as significant results from immune simulation demonstrated the efficacy of the vaccine in the immune cell interaction during the innate and adaptive immune responses immune simulation. In conclusion, the outcome suggested that the newly developed vaccine will be a promising candidate for controlling and providing definitive preventive measures against scrub typhus if further investigation is conducted experimentally.



URL: <https://www.frontiersin.org/journals/immunology/articles/10.3389/fimmu.2025.1513245/full>





SCHOLARLY PUBLICATIONS
School of Biotechnology
KIIT Deemed to be University

Journal Name: Materials Chemistry and Physics

IF: 4.7

Title: Deciphering biosurfactant-salt interaction and its influence on biosurfactant activity in muga silk fibroin extraction

Author: Biswal B.; Das M.; Das D.; Prusty D.; Dan A.K.

Details: Volume 344, October 2025

Abstract: Degumming of silk cocoons is the initial technique employed to separate two silk proteins (fibroin and sericin), which further leads to the formulation of diverse silk-based biomaterials for biomedical applications. In this study, a novel approach has been implemented in a mixed system. Specifically, this paper emphasizes the impact of degumming on fibroin fiber, which was carried out using varying concentrations of sodium carbonate (Na_2CO_3) and crude biosurfactant extracted from *Acacia concinna* (Willd.) Dc. The study investigated the effectiveness of the degumming process under a specific concentration of Na_2CO_3 with crude biosurfactant extract, examining the influence of reaction time, temperature, and mixed reagent concentration. The results of the degumming process show that approximately 24.8 % degumming occurred when using 0.012 g/mL of biosurfactant extract with 3×10^{-4} g/mL of Na_2CO_3 as degumming reagents. Furthermore, SEM, XRD, TGA, and mechanical strength analyses suggest that the quality of fibers extracted using the crude biosurfactant (BSE) and Na_2CO_3 mixture in the degumming process yielded significant results. This innovative approach of degumming can extract the silk fibroin from the cocoons in the fastest and most effective way. Moreover, this strategy may significantly diminish the harmful contamination of sericin and degumming chemicals in the effluent.



URL: <https://www.sciencedirect.com/science/article/pii/S0254058425008107?via%3Dihub>





SCHOLARLY PUBLICATIONS
School of Biotechnology
KIIT Deemed to be University

Journal Name: Bioorganic Chemistry


IF: 4.7

Title: Design, synthesis and biological evaluation of 2-phenylquinoxaline carbonyl piperazine derivatives as novel FASN inhibitors with anticancer activity

Author: Singh S.; Paul S.; Martins F.G.; Sousa S.F.; Kundu C.N.; Karthikeyan C.; Moorthy N.S.H.N.

Details: Volume 163, August 2025

Abstract: Overexpression of fatty acid synthase (FASN) has been linked to the advancement of various cancers. FASN caters to the increased demand for lipids within tumor cells, facilitating tumor growth and progression, making it an attractive target for anticancer drug discovery. Herein we report a novel series of 2-phenylquinoxaline-6-carboxylic acid derivatives as novel potent FASN inhibitors with anticancer potential. Structure-activity relationship analysis demonstrated that all the synthesized compounds showed potent and selective cytotoxicity against the three cancer cell lines evaluated with IC_{50} values less than 10 μ M. QNX-10 was identified as a promising lead molecule as it elicited potent FASN inhibition and selective cytotoxicity against the colorectal (HCT-116, Caco-2 cell lines) and breast cancer (MCF-7 cell line). Notably, QNX-10 induces apoptosis and cell cycle arrest at S-phase in HCT-116 cells in a dose-dependent manner. Western blot analysis indicated that QNX-10 inhibits FASN and promotes apoptosis in HCT-116 cells by upregulating pro-apoptotic protein Bax and downregulating anti-apoptotic protein Bcl-xL. Molecular docking and MD simulation studies with QNX-10 revealed the binding mode of the compound to the KR domain of FASN. Taken together, the study establishes compound QNX-10 to be a promising lead candidate for the development of anticancer therapeutics targeting the FASN enzyme.



URL: <https://www.sciencedirect.com/science/article/abs/pii/S0045206825005772?via%3Dihub>





SCHOLARLY PUBLICATIONS School of Biotechnology KIIT Deemed to be University

Journal Name: Materials Advances

IF: 4.7

Title: Bactericidal activity of ZnO nanoparticles-anti-TB drug combination towards the H37Rv strain and multidrug-resistant isolates of Mycobacterium tuberculosis via SufB splicing inhibition

Author: Ojha D.K.; Mehra A.; Rout S.S.; Giri S.; Nayak S.

Details: Issue 15, 2025

Abstract: Tuberculosis (TB) remains a significant global health threat, claiming millions of lives annually. Despite advancements in treatment, the emergence of drug-resistant strains has hindered effective TB control. The current management of TB involves prolonged treatment duration with severe side effects, leading to poor patient compliance. However, the mycobactericidal potential of nanoparticles towards drug-resistant TB is not confirmed yet. This work explores the bactericidal potential of zinc oxide nanoparticles (ZnONPs, 40 nm) in managing both drug-sensitive and drug-resistant TB in combination with anti-TB drugs. It was found that ZnONPs inhibit the generation of active SufB protein via splicing inhibition, an essential event for Mycobacterium tuberculosis (Mtb) survival. While TEM and UV-visible spectroscopy identified NPs-protein interaction, SEM visualised extensive membrane damage in H37Rv and multidrug-resistant (MDR) Mtb cells. Alamar blue assay and the spread plate method detected minimum inhibitory concentration and minimum bactericidal concentration of ZnONPs towards the H37Rv strain and MDR Mtb isolates. In vitro studies identified a combination with ZnONPs that reduced effective doses for anti-TB drugs towards H37Rv and MDR Mtb isolates. A similar drug combination attenuated the mycobacterial load and inflammation in the spleen and lungs and protected against Mtb induced splenomegaly in infected mice. Thus, ZnONPs can be used as a potent additive in the anti-TB regimen to manage drug-susceptible and drug-resistant TB, addressing challenges such as prolonged therapy, drug toxicity and poor patient compliance.



URL: <https://pubs.rsc.org/en/content/articlelanding/2025/ma/d4ma01224k>





SCHOLARLY PUBLICATIONS School of Biotechnology KIIT Deemed to be University

Journal Name: International Journal of Environmental Research and Public Health **IF:** 4.6

Title: Microplastic Pollution: A Global Environmental Crisis Impacting Marine Life, Human Health, and Potential Innovative Sustainable Solutions

Author: Karak P.; Parveen A.; Modak A.; Adhikari A.; Chakraborty S.

Details: Vol 22, Issue 6, Jun 2025

Abstract: Pollution, especially plastic pollution, presents a serious worldwide danger to essential environmental resources. Microplastics are tiny plastic fragments varying in size from 50 μm to 5 mm. The primary aim of this article is to develop an extensive review grounded in the latest data accessible until 2024, adhering to PRISMA guidelines. A total of 329 data points were collected and 297 of those were removed through filtering, leaving 32 articles for the study, and taking into account the complete evolution of all the publications. This study seeks to enhance public awareness and knowledge among researchers about the harmful effects of plastic pollution on the environment and society by identifying its sources and consequences for humans and ecosystems. A detailed analysis of the sources of microplastics in the oceans and their detrimental effects on marine organisms is presented. This research additionally explores the transport of microplastics through various environmental pathways, including water and air. Aquatic species ingest microplastics, which subsequently transfer up the food chain, including humans, and these risks are discussed. Microplastics may increase the production of reactive oxygen species (ROS), leading to DNA and cellular damage, oxidative stress, alterations in gene expression, and decreased cell viability. Developing clear and effective guidelines and regulations is crucial for addressing the adverse issues related to microplastics. All participants in the policymaking and implementation of these guidelines must understand their roles and responsibilities.



URL: <https://www.mdpi.com/1660-4601/22/6/889>





SCHOLARLY PUBLICATIONS School of Biotechnology KIIT Deemed to be University

Journal Name: Spectrochimica Acta - Part A: Molecular and Biomolecular Spectroscopy

IF:4.6

Title: Spectroscopic investigation of hydrogen bond network stability and microplastic leaching in ethanol-based potentised medicines at extreme dilutions during prolonged plastic storage

Author: Chakraborty S.; Ghosh K.; Biswas S.; Roy Chaudhuri C.; Roy Chowdhury A.; Chakravarty R.; Nayak D.; Kaushik S.; Barui A.; Kundu S.

Details: Vol. 343, Dec 2025

Abstract: The quality and efficacy of pharmaceutical products stored under proper conditions are critical. This study examined the effects of long-term plastic storage on extremely diluted ethanol-based potentised (EP) medicines using advanced spectroscopic techniques. Four medicines, Arnica montana, Rhus toxicodendron, Conium maculatum, and Belladonna, at ultra-high (200C, 1 M) and moderate-high (30C, 200C) potencies, were stored in glass and plastic containers for one month. Glass-stored medicines showed increased antioxidant activity and zeta potential with higher potency, while plastic-stored samples showed a decreasing trend. Conductivity was inversely correlated with zeta potential, with glass-stored medicines showing a $\sim 41.91\%$ reduction, while plastic-stored samples showed a $\sim 36.29\%$ increase. Mid-IR spectra revealed a blue shift ($\sim 4\text{--}14\text{ cm}^{-1}$) in O–H stretching and a red shift ($\sim 2\text{--}3\text{ cm}^{-1}$) in H–O–H bending for glass-stored medicines, showing weaker inter-molecular H-bonds at higher potencies. In contrast, plastic-stored medicines showed opposite shifts ($\sim 2\text{--}17\text{ cm}^{-1}$), implying more constrained H-bonding due to carbonyl-water interaction in presence of microplastics, disrupting the native ethanol-water H-Bond network. Far-IR spectra showed an enthalpic gain ($\sim 45.34\%$) in glass-stored medicines, while plastic-stored samples showed an enthalpic loss ($\sim 56.60\%$), confirming structural destabilisation of native water-network due to microplastic leaching. Our findings show that plastic containers compromised the efficacy of studied medicines by altering H-bond network stability and electrical properties. Further studies on different plastic grades and storage durations are needed to validate these findings and explore cost-effective alternatives for long-term storage of such medicines.



URL: <https://www.sciencedirect.com/science/article/abs/pii/S1386142525009229?via%3Dihub>





SCHOLARLY PUBLICATIONS
School of Biotechnology
KIIT Deemed to be University

Journal Name: Inorganic Chemistry Communications


IF: 4.4

Title: Bio-synthesized cerium oxide nano-particles for efficient adsorption of fluoride ion from aqueous solution

Author: Hansdah S.; Das J.; Mandal D.; Parhi P.K.

Details: Volume 178, Part 2, August 2025, 114583

Abstract: Cerium oxide (CeO_2) nanoparticles (NPs) were successfully synthesized via a facile biosynthesis route, utilizing a cost-effective and environmentally friendly bio-extract derived from fenugreek seeds, for the effective removal of fluoride ions from contaminated water. The synthesized CeO_2 NPs were characterized using UV, FTIR, XRD and SEM to determine the phase, purity, and size, revealing an average size of approximately 17.4 nm. The fluoride sorption behavior and mechanism on to the ceria NPs phase was confirmed through kinetics, isotherm, and thermodynamics investigations. The sorption affinity of fluoride ion with ceria NPs appears to be effective at solution pH 2.0. The fluoride sorption rate onto the nano-ceria phase was observed to follow a pseudo-second order kinetic model along with intra-particle diffusion. Among the three sorption adopted isotherm models; Langmuir, Freundlich and Temkin, the fluoride adsorption behavior revealed to follow Langmuir Isotherm with high regression coefficient value ($R^2 = 0.993$). The maximum fluoride adsorption loading capacity of CeO_2 NPs was determined to be 123.6 mg/g at the temp. 293 K. The fluoride ion adsorption with CeO_2 NPs is observed to be spontaneous and endothermic in nature which was strongly supported by the thermodynamics results; $\Delta H = 13.789$ KJ/mol, and $\Delta S = +62.799$. The phase transformation due to the sorption of fluoride onto the homogeneous ceria phase was ascertained isotherm results and EDX analysis resulted of fluoride ion loaded Ceria NPs.



URL: <https://www.sciencedirect.com/science/article/pii/S1387700325006999?via%3Dihub>





SCHOLARLY PUBLICATIONS School of Biotechnology KIIT Deemed to be University

Journal Name: Frontiers in Pharmacology


IF: 4.4

Title: Editorial: Emerging and reemerging neglected tropical diseases: epidemiology, transmission, mitigation, and vaccines and chemotherapy advancements

Author: Mohapatra R.K.; Mishra S.; Kandi V.; Sirka C.S.; Tuglo L.S.

Details: Volume 16, 2025, Article number 1545801

Abstract: Neglected tropical diseases (NTDs) are infectious diseases caused by bacteria, viruses, fungi and parasites, including ectoparasites like mites and fleas ([Mohapatra et al., 2024](#); [Kutikuppala et al., 2023](#); [WHO, 2024](#)). NTDs are an ongoing challenge to global public healthcare and community health. The main reason why such diseases remain collectively neglected is that they are considered “diseases of the poor”, primarily people in low- and middle-income countries (LMICs) with modest purchasing capacity. As a result of this, the commercial diagnostic, therapeutic and prophylactic efforts by the pharmaceutical companies are only skeletal as they do not envision a profitable market. Thus, although treatable, these diseases ultimately manifest as terminal diseases of the have-nots. The World Health Organisation (WHO) has compiled a list of the world’s most prevalent NTDs and updates it from time to time. Currently, there are 21 aetiologically, epidemiologically and clinically unique diseases (or groups of diseases) listed as NTDs by the WHO ([Malecela and Ducker, 2021](#)). The WHO has devised public health strategies and proposed a roadmap to eliminate NTDs by 2030 ([WHO, 2021](#)). However, as the majority of the NTD-affected subjects live in financially constrained third-world countries, this seems to be an uphill task and achieving it is by no means easy. Lack of NTD-related awareness and limited diagnostic resources in these regions severely affect their foolproof identification. This is evidenced by the constantly increasing number of diseases that fit into the WHO criteria for NTDs. Various NTDs re-emerged in the wake of the coronavirus disease 2019 (COVID-19) pandemic by the severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2), which hit the healthcare infrastructure hard worldwide and exposed its underbelly.



URL: <https://www.frontiersin.org/journals/pharmacology/articles/10.3389/fphar.2025.1545801/full>





SCHOLARLY PUBLICATIONS
School of Biotechnology
KIIT Deemed to be University

Journal Name: Journal of Molecular Structure


IF: 4.0

Title: Antimicrobial activities of novel substituted spiropyrrolidine based heterocycles synthesized by multicomponent reaction against *Bacillus subtilis* and *Pseudomonas aeruginosa*

Author: Suresh Babu A.R.; Rani S.; Singh S.P.; Khera A.; Alajangi H.K.; Parandaman S.; Raj A.R.N.; Gavaskar D.; Gartia J.; Pandey A.; Yadav V.K.; Singh G.; Barnwal R.P.

Details: Volume 1339, 5 September 2025, Article number 142373

Abstract: The growing resistance of bacteria to antimicrobial agents has intensified the need for novel strategies to combat bacterial infections, particularly those associated with biofilm formation. Biofilms enhance bacterial resilience against hostile environments, immune responses, and antimicrobial treatments. The ability of biofilms to influence bacterial pathogenesis underscores the critical need for new antibacterial agents with anti-biofilm activity. This research aims to synthesize cost-effective, structurally diverse, chemical compounds with the biological significance of disrupting biofilm formation. Here, we report a facile sequential reaction for one-pot, four-component synthesis of spiropyrrolidine heterocycles with 1,3-dipolar cycloaddition of azomethine ylide. The multicomponent reaction (MCR) provides high yield and regioselectivity of the desired product, under mild reaction conditions. Preliminary screening for these novel compounds involves biofilm assays, which assess the developmental processes of biofilms, providing insights into the compounds' biological potential. Subsequent in vitro experiments assessed their antibacterial potential against *B. subtilis* and *P. aeruginosa* using the minimum inhibitory concentration (MIC) assay. A cell culture assay evaluated toxicity of these compounds in MDA-MB-231 cell lines. All these investigations cumulatively highlight the potential of these molecules as antibacterial agents for *B. subtilis* and *P. aeruginosa*.



URL: <https://www.sciencedirect.com/science/article/pii/S0022286025010531?via%3Dihub>





SCHOLARLY PUBLICATIONS School of Biotechnology KIIT Deemed to be University

Journal Name: Scientific Reports

IF: 3.9

Title: Novel endophytic actinomycetes species *Streptomyces panacea* of *Panax sokpayensis* produce antimicrobial compounds against multidrug resistant *Staphylococcus aureus*

Author: Rai S.; Singh L.S.; Liriina K.; Jeyaram K.; Parija T.; Sahoo D.

Details: Volume 15, Issue 1, December 2025, Article number 19863

Abstract: Endophytic actinomycetes of medicinal plants have recently been in focus for developing novel antimicrobial compounds to combat multidrug-resistant pathogens. In this study, we isolated and characterised endophytic actinomycetes of *Panax sokpayensis* rhizome traditionally used as medicine in Sikkim-Himalayan region and assessed their antimicrobial activity against multidrug-resistant (MDR) clinical isolates of *Staphylococcus aureus*. *Saccharopolyspora* dominated as the endophytic actinomycetes of *P. sokpayensis* rhizome. However, a novel actinomycete strain PSRA5^T belongs to the genus *Streptomyces*, with the highest genome sequence similarity of 91.54% with its closest relative *Streptomyces niveus* NCIMB 11891 has shown an effective inhibition of six clinical isolates of MDR *S. aureus* during disc diffusion assay. Further comparative analysis of cellular fatty acids composition and phenotypic and biochemical characteristics of strain PSRA5^T with its phylogenetically closely related strain of *S. niveus*, classified as representing a novel species of the genus *Streptomyces*, for which the name *Streptomyces panacea* sp. nov. is proposed here with type strain PSRA5^T (= MCC5238^T). The minimum inhibition concentration of ethyl acetate crude extract of PSRA5^T culture supernatant against MDR *S. aureus* isolates was 5.5 to 13.5 µg/mL. Further correlation between biosynthetic gene clusters identified by genome search with LC-MS analysis-based chemical profiling of PSRA5^T culture extract and antibacterial activity of the representative compounds detected several compounds of aminoglycosides and polyketides with antimicrobial activity against MDR *S. aureus* isolates.



URL: <https://www.nature.com/articles/s41598-025-05333-1>





SCHOLARLY PUBLICATIONS School of Biotechnology KIIT Deemed to be University

Journal Name: Langmuir


IF: 3.9

Title: Luminescent Carbon Dot-Graphene Oxide Epoxy Coatings: Enhanced Corrosion Protection and Detection

Author: Daksha D.; Rout T.K.; Bhagat A.N.; Satpathy B.K.; Mishra S.; Panigrahi H.

Details: Vol 41, Issue 28, July 2025

Abstract: A new hybrid material-based approach leveraging luminescent hybrid carbon dots (CDs)-functionalized graphene oxide (CDs-f-GO) integrated into epoxy coatings has been developed for enhanced anticorrosion protection and real-time corrosion detection. This study introduces, for the first time, a one-step process for synthesizing CDs-functionalized GO, which has been thoroughly characterized by using Fourier transform infrared spectroscopy, UV-visible spectroscopy, and fluorescence spectroscopy, confirming successful functionalization. UV-visible spectra revealed a characteristic absorption peak at 335 nm for CDs-f-GO, while fluorescence spectroscopy demonstrated excitation-dependent fluorescence with a red shift from 445 to 472 nm, distinguishing it from nonfluorescent GO. Electrochemical characterization, including electrochemical impedance spectroscopy and Tafel analysis, confirmed the superior anticorrosion performance of epoxy-CDs-f-GO coatings. Corrosion resistance was evaluated over time in a 3.5 wt % NaCl solution, revealing significantly higher charge transfer resistance and lower corrosion current density for epoxy-CDs-f-GO compared to neat epoxy and other composite coatings. The hybrid material exhibited the highest corrosion protection on both days 1 and 7, maintaining minimal degradation over time. The salt spray test on scratched coatings further validated the effectiveness of epoxy-CDs-f-GO, as minimal corrosion blistering was observed, indicating strong resistance against electrolyte penetration. Additionally, the developed coating demonstrated dual functionality by enabling real-time failure detection under UV light. Prominent carving marks were visible on the epoxy-CDs-f-GO coating, suggesting its potential for early detection of cracks or defects in steel structures. The combination of enhanced corrosion resistance and proactive maintenance capability makes this material a promising candidate for industrial applications, particularly in marine and other highly corrosive environments.



URL: <https://pubs.acs.org/doi/10.1021/acs.langmuir.5c00875>





SCHOLARLY PUBLICATIONS
School of Biotechnology
KIIT Deemed to be University

Journal Name: Plant Signaling and Behavior

IF: 3.6

Title: Arbuscular mycorrhizal fungi—a natural tool to impart abiotic stress tolerance in plants

Author: Samanta I.; Ghosh K.; Saikia R.; Savita; Maity P.J.; Chowdhary G.

Details: Volume 20, Issue 1, July 2025

Abstract: Arbuscular mycorrhizal fungi (AMF) are crucial components of the soil microbiomes that establish symbiotic associations with most terrestrial plants. The review summarizes the basic mechanisms behind the plant-AMF symbiosis, the genes involved in the fungal and their plant counterparts, novel biomolecules and growth regulators, leading to probable signal transduction pathways. It also focuses on the involvement of lipids and strigolactones in establishing AMF-plant symbiosis. Herein, we further emphasize the role played by these AMF in enhancing plant resistance to various abiotic stresses while giving a broad outline of current research practices and attempting to dissect the mechanism behind the AMF-mediated abiotic stress signal transduction. Discussion on the mechanisms behind this stress reduction involving AMF will be valuable for the researchers, agronomists, and environmentalists involved in sustainable agriculture. Water scarcity, salinity, heavy metals, and extreme temperatures are the primary abiotic stresses that pose serious challenges to agricultural sustainability and ecosystem functioning. Conventional responses to such pressures typically rely on genetic modifications as well as chemical treatments, which could be expensive and detrimental to the environment. However, these AM fungi act in an alternative way that is natural and cost-effective too, leading to healthy plants with resilience toward stress through symbiosis, leading to the fulfillment of the United Nations Sustainable Development Goal (UNSDG) 2 of zero hunger.



URL: <https://www.tandfonline.com/doi/full/10.1080/15592324.2025.2525843>

